

Low-temperature thermodynamics of one-dimensional alternating-spin Heisenberg ferromagnets

Shoji Yamamoto and Hiromitsu Hori

Division of Physics, Hokkaido University, Sapporo 060-0810, Japan

(Dated: 27 April 2005)

Motivated by a novel bimetallic chain compound in which alternating magnetic centers are ferromagnetically coupled, we investigate thermodynamic properties of one-dimensional spin- (S, s) Heisenberg ferromagnets both numerically and analytically. On the one hand, quantum Monte Carlo calculations illuminate the overall thermal behavior. The specific heat may exhibit a double-peaked structure at intermediate temperatures for $S \gtrsim 3s$ in general. On the other hand, a modified spin-wave theory precisely describes the low-temperature properties. Expanding the specific heat and the magnetic susceptibility, we reveal an analogy and a contrast between mixed-spin ferromagnets and ferrimagnets.

PACS numbers: 75.10.Jm, 75.40.Cx, 75.30.Ds, 75.40.Mg

I. INTRODUCTION

Much effort has been devoted to designing molecular systems ordering ferromagnetically.¹ One possible approach consists of assembling molecular bricks so as to obtain a low-dimensional system with a nonzero resultant spin in the ground state and then coupling the chains or the layers again in a ferromagnetic fashion. Numerous heterospin chain compounds have been synthesized along this line. Gleizes and Verdager² made an attempt to alternate two types of metal ion along one crystallographic axis with antiferromagnetic intrachain interaction and obtained a pioneering example of quasi-one-dimensional ferrimagnets, of formula $\text{MnACu}(\text{dto})_2(\text{H}_2\text{O})_3 \cdot 4.5\text{H}_2\text{O}$ ($A = \text{Ni, Cu}$; $\text{dto} = \text{dithiooxalato} = \text{S}_2\text{C}_2\text{O}_2$). Kahn *et al.*³ synthesized another series of bimetallic chain compounds $\text{ACu}(\text{pbaOH})(\text{H}_2\text{O})_3 \cdot n\text{H}_2\text{O}$ ($A = \text{Fe, Co, Ni, Cu}$; $\text{pbaOH} = 2\text{-hydroxy-1, 3-propylenebis(oxamate)} = \text{C}_7\text{H}_6\text{N}_2\text{O}_7$), one of which indeed attained the three-dimensional ferromagnetic order at low temperatures.⁴ Caneschi *et al.*⁵ took an alternative strategy of bringing into interaction metal ions and stable organic radicals. This idea was developed toward polymeric chain compounds.⁶ The wide variety of chemical explorations stimulated the physical interest in mixed-spin chains.^{7,8,9,10,11,12,13,14,15,16}

Most of the thus-far synthesized heterospin systems are characterized as ferrimagnets. Ferromagnetic intrachain coupling is observed in few cases. In such circumstances, $\text{MnNi}(\text{NO}_2)_4(\text{en})_2$ ($\text{en} = \text{ethylenediamine} = \text{C}_2\text{H}_8\text{N}_2$),¹⁷ proved to be a quasi-one-dimensional mixed-spin ferromagnet¹⁸ and caused renewed interest in mixed-spin chains. Gillon *et al.*¹⁹ calculated the spin density distribution by means of the density functional theory and quantitatively visualized the ferromagnetic nature of the Mn(II)-Ni(II) interaction. Fukushima *et al.*²⁰ performed high-temperature series expansion of the thermal quantities and argued the magnetic structure including single-ion anisotropy and interchain exchange coupling. Now an increasing number of chemists and physicists are taking interest in heterospin ferromagnets.^{21,22}

Alternating-spin chains possess elementary excitations of dual aspect. In the case of antiferromagnetic coupling, the acoustic excitations reduce the ground-state magnetization and are thus of ferromagnetic nature, while the optical excitations enhance the ground-state magnetization and are thus of antiferromagnetic nature. In the case of ferromagnetic coupling, on the other hand, both excitations are of ferromagnetic character. Therefore, the Schottky-type peak of the specific heat and the minimum of the susceptibility-temperature product, which are both ferrimagnetic features, are absent from mixed-spin ferromagnets. Nevertheless, mixed-spin ferromagnets and ferrimagnets behave similarly at low temperatures, which is the goal of this paper. Employing a quantum Monte Carlo method²³ and a modified spin-wave theory,²⁴ we investigate thermodynamics of one-dimensional alternating-spin Heisenberg ferromagnets with particular emphasis on the intrinsic low-temperature properties.

II. MODIFIED SPIN-WAVE SCHEME

We consider two kinds of spins S and s ($S > s$) alternating on a ring with ferromagnetic exchange coupling between nearest neighbors, as described by the Hamiltonian

$$\mathcal{H} = -J \sum_{n=1}^N (\mathbf{S}_n \cdot \mathbf{s}_n + \mathbf{s}_n \cdot \mathbf{S}_{n+1}). \quad (2.1)$$

Even in one dimension, the conventional spin-wave theory^{25,26,27} gives a fine piece of information on the ground-state correlation.^{28,29} As for the thermal quantities, however, the low-temperature series expansion within the conventional scheme³⁰ only reproduces the leading term of the specific heat and nothing correct for the magnetic susceptibility.³¹ Then Takahashi³² modified the spin-wave formalism, imposing a constraint on the magnetization, and obtained an excellent description of the low-temperature thermodynamics of low-dimensional

ferromagnets. We develop the modified scheme for mixed-spin ferromagnets.

In order to describe the spin deviation in each sublattice, bosonic operators are introduced as

$$\begin{aligned} S_n^+ &= \sqrt{2S - a_n^\dagger a_n} a_n, \quad S_n^z = S - a_n^\dagger a_n, \\ s_n^+ &= \sqrt{2s - b_n^\dagger b_n} b_n, \quad s_n^z = s - b_n^\dagger b_n, \end{aligned} \quad (2.2)$$

where we regard S and s as quantities of the same order. The bosonic Hamiltonian reads

$$\mathcal{H} = E_2 + \mathcal{H}_1 + \mathcal{H}_0 + O(S^{-1}), \quad (2.3)$$

where $E_2 = -2SsJN$ is the classical ground-state energy and $\mathcal{H}_i$ is the $O(S^i)$ quantum correction to it. We consider first diagonalizing $\mathcal{H}_1$ and then taking $\mathcal{H}_0$ into calculation perturbationally.³³ Via the transformation

$$\begin{aligned} a_n^\dagger &= \frac{1}{\sqrt{N}} \sum_k e^{-ik(n-1/4)} (\alpha_k^\dagger \cos \theta_k - \beta_k^\dagger \sin \theta_k), \\ b_n^\dagger &= \frac{1}{\sqrt{N}} \sum_k e^{-ik(n+1/4)} (\alpha_k^\dagger \sin \theta_k + \beta_k^\dagger \cos \theta_k), \end{aligned} \quad (2.4)$$

with $\tan(2\theta_k) = 2\sqrt{Ss} \cos(k/2)/(S-s)$, we obtain

$$\mathcal{H}_1 = J \sum_k (\omega_k^- \alpha_k^\dagger \alpha_k + \omega_k^+ \beta_k^\dagger \beta_k). \quad (2.5)$$

Here the acoustic (ω_k^-) and optical (ω_k^+) dispersion relations are given by

$$\begin{aligned} \omega_k^\pm &= S + s \pm \sqrt{(S-s)^2 + 4Ss \cos^2(k/2)} \\ &\equiv S + s \pm \omega_k, \end{aligned} \quad (2.6)$$

and plotted in Fig. 1.

Now we proceed to the modified spin-wave scheme in an attempt to avoid thermal divergence of the number of bosons. At finite temperatures, we replace $\alpha_k^\dagger \alpha_k$ and $\beta_k^\dagger \beta_k$ by $\bar{n}_k^\mp \equiv \sum_{n^-, n^+} n^\mp P_k(n^-, n^+)$, where $P_k(n^-, n^+)$ is the probability of n^- acoustic and n^+ optical spin waves appearing in the k -momentum state, and minimize the up-to- $O(S^1)$ free energy

$$\begin{aligned} \mathcal{F} &= E_2 + J \sum_k \sum_{\sigma=\pm} \omega_k^\sigma \bar{n}_k^\sigma \\ &+ k_B T \sum_k \sum_{n^-, n^+} P_k(n^-, n^+) \ln P_k(n^-, n^+), \end{aligned} \quad (2.7)$$

with respect to $P_k(n^-, n^+)$'s under the condition of zero magnetization

$$(S+s)N - \sum_k \sum_{\sigma=\pm} \bar{n}_k^\sigma = 0, \quad (2.8)$$

together with the trivial constraints $\sum_{n^-, n^+} P_k(n^-, n^+) = 1$. Up to $O(S^1)$, the magnetic susceptibility and the internal energy at thermal

equilibrium are expressed as

$$\chi = \frac{(g\mu_B)^2}{3k_B T} \sum_k \sum_{\sigma=\pm} \bar{n}_k^\sigma (\bar{n}_k^\sigma + 1), \quad (2.9)$$

$$E = E_2 + J \sum_k \sum_{\sigma=\pm} \omega_k^\sigma \bar{n}_k^\sigma, \quad (2.10)$$

with $\bar{n}_k^\pm = [e^{(J\omega_k^\pm - \mu)/k_B T} - 1]^{-1}$, where the g factors of the spins S and s are both set equal to g and the Lagrange multiplier μ is determined through the condition (2.8). The specific heat is calculated by numerically differentiating the internal energy. The perturbational correction of $O(S^0)$ reads

$$\begin{aligned} \langle H_0 \rangle &\equiv \text{Tr}[\mathcal{H}_0 e^{-\mathcal{H}_1/k_B T}] / \text{Tr}[e^{-\mathcal{H}_1/k_B T}] \\ &= \frac{JN}{2} \left[\sqrt{\frac{S}{s}} (\Gamma_1 - \Gamma_2) \Gamma_3 + \sqrt{\frac{s}{S}} (\Gamma_1 + \Gamma_2) \Gamma_3 \right. \\ &\quad \left. - \Gamma_1^2 + \Gamma_2^2 - \Gamma_3^2 \right], \end{aligned} \quad (2.11)$$

with

$$\begin{aligned} \Gamma_1 &= \frac{1}{N} \sum_k (\bar{n}_k^- + \bar{n}_k^+) = S + s, \\ \Gamma_2 &= \frac{1}{N} \sum_k \frac{S-s}{\omega_k} (\bar{n}_k^- - \bar{n}_k^+), \\ \Gamma_3 &= \frac{1}{N} \sum_k \frac{2\sqrt{Ss}}{\omega_k} \cos^2 \frac{k}{2} (\bar{n}_k^- - \bar{n}_k^+), \end{aligned} \quad (2.12)$$

where we keep μ unchanged. Indeed μ may be modified so as to minimize the up-to- $O(S^0)$ free energy, but such a procedure, which is much more complicated, has no effect on the low-temperature leading behavior.³⁰

III. NUMERICAL CALCULATIONS

Quantum Monte Carlo calculations are presented at $N = 32$ (64 spins). In the case of $(S, s) = (1, \frac{1}{2})$, we have carried out preliminary calculations at $N = 24, 32, 40$. Any quantity divided by N does not vary with N beyond its statistical error in the temperature range to show. A few million Monte Carlo steps are spent on low-temperature calculations, while less than a half million steps on high-temperature calculations. The numerical precision in the final results is two to three digits.

In Fig. 2 we compare modified-spin-wave and quantum Monte Carlo calculations of the magnetic susceptibility. They are in excellent agreement over the whole temperature range. The analytic calculation reproduces the paramagnetic susceptibility $[S(S+1)+s(s+1)]N(g\mu_B)^2/3k_B T$ at high temperatures and reveals the T^{-2} -diverging behavior at low temperatures, which is later discussed in more detail. The susceptibility-temperature product still monotonically decreases with increasing temperature in contrast with the ferrimagnetic behavior.³⁴

In Fig. 3 we compare the modified spin-wave and quantum Monte Carlo calculations of the specific heat. The agreement between them is somewhat poorer than that found for the susceptibility, but the $T^{1/2}$ -initial behavior at low temperatures and the spin-dependent peak structure at intermediate temperatures are well reproduced by the analytic calculation. The mid-temperature structure of the specific heat may be regarded as a function of the acoustic excitation band width $W^- = 2sJ$ and the optical excitation gap $\Delta = 2SJ$ (see Fig. 1). The heat capacity attributable to the acoustic excitations and that to the optical excitations may be separable when $W^- \ll \Delta$. Observing further calculations in Fig. 4, we are convinced that the double-peaked structure may appear for $S \gtrsim 3s$, including practical cases $(S, s) = (\frac{5}{2}, \frac{1}{2}), (2, \frac{1}{2}), (\frac{3}{2}, \frac{1}{2})$. Mn(II)Cu(II), Fe(II)Cu(II), and Co(II)Cu(II) chain compounds³ have indeed been synthesized so far, but they all exhibit antiferromagnetic intrachain interaction. The double-peaked structure is much more pronounced for ferromagnetic intrachain interaction.^{20,35} We expect an increased effort to design ferromagnetic exchange coupling between alternating metal ions.

IV. ANALYTICAL CALCULATIONS

In order to elucidate the low-temperature thermal behavior, we define the state density function

$$w^\pm(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \delta(x - \omega_k^\pm) dk. \quad (4.1)$$

Here we are interested in the gapless acoustic branch and expand $w^-(x)$ for small x as

$$w^-(x) = \frac{1}{\pi} \sqrt{\frac{S+s}{2Ss}} \sum_{n=0}^{\infty} c_n^- x^n. \quad (4.2)$$

A few leading coefficients are given as

$$\begin{aligned} c_0^- &= 1, \\ c_1^- &= \frac{(S-s)^2 + Ss}{4Ss(S+s)}, \\ c_2^- &= \frac{(3S^2 - 4Ss + 3s^2)(S+s)^2 - 5S^2s^2}{32S^2s^2(S+s)^2}. \end{aligned} \quad (4.3)$$

Applying Eq. (4.2) and neglecting the optical excitations $\bar{n}_k^+$, Eq. (2.8) reads

$$\begin{aligned} v^{1/2} &= \frac{1}{\pi \sqrt{2Ss(S+s)}} \sum_{n=0}^{\infty} c_n^- t^{n+1/2} \\ &\times \Gamma\left(n + \frac{1}{2}\right) \left[\Gamma\left(\frac{1}{2} - n\right) v^n \right. \\ &\left. + \sum_{m=0}^{\infty} \frac{(-1)^m}{m!} \zeta\left(n - m + \frac{1}{2}\right) v^{m+1/2} \right], \end{aligned} \quad (4.4)$$

where $v = -\mu/k_B T$, $t = k_B T/J$, and $\zeta(n)$ is Riemann's zeta function. Solving this equation iteratively, we obtain the low-temperature expansion of the Lagrange multiplier as

$$\begin{aligned} v^{1/2} &= \frac{c_0^- \Gamma(1/2)}{\pi \sqrt{2Ss(S+s)}} \Gamma\left(\frac{1}{2}\right) t^{1/2} + \left[\frac{c_0^- \Gamma(1/2)}{\pi \sqrt{2Ss(S+s)}} \right]^2 \\ &\times \Gamma\left(\frac{1}{2}\right) \zeta\left(\frac{1}{2}\right) t + \left[\frac{c_0^- \Gamma(1/2)}{\pi \sqrt{2Ss(S+s)}} \right]^3 \\ &\times \Gamma\left(\frac{1}{2}\right) \left[\zeta\left(\frac{1}{2}\right) \right]^2 t^{3/2} + O(t^2). \end{aligned} \quad (4.5)$$

Since the magnetic susceptibility and the internal energy read

$$\begin{aligned} \frac{\chi k_B T}{N(g\mu_B)^2} &= \frac{1}{3\pi} \sqrt{\frac{S+s}{2Ss}} \sum_{n=0}^{\infty} c_n^- t^{n+1/2} \\ &\times \Gamma\left(n + \frac{1}{2}\right) \left[\Gamma\left(\frac{3}{2} - n\right) v^{n-3/2} \right. \\ &\left. + \sum_{m=0}^{\infty} \frac{(-1)^m}{m!} \zeta\left(n - m - \frac{1}{2}\right) v^m \right], \end{aligned} \quad (4.6)$$

$$\begin{aligned} \frac{E - E_2}{NJ} &= \frac{1}{\pi} \sqrt{\frac{S+s}{2Ss}} \sum_{n=0}^{\infty} c_n^- t^{n+3/2} \\ &\times \Gamma\left(n + \frac{3}{2}\right) \left[\Gamma\left(-\frac{1}{2} - n\right) v^{n+1/2} \right. \\ &\left. + \sum_{m=0}^{\infty} \frac{(-1)^m}{m!} \zeta\left(n - m + \frac{3}{2}\right) v^m \right], \end{aligned} \quad (4.7)$$

the susceptibility and the specific heat are expanded as

$$\begin{aligned} \frac{\chi J}{N(g\mu_B)^2} &= \frac{1}{Ss} \left\{ \frac{\tilde{t}^{-2}}{3} - \frac{\zeta(1/2)}{\sqrt{2\pi}} \tilde{t}^{-3/2} \right. \\ &\left. + \left[\frac{\zeta(1/2)}{\sqrt{2\pi}} \right]^2 \tilde{t}^{-1} \right\} + O(\tilde{t}^{-1/2}), \end{aligned} \quad (4.8)$$

$$\begin{aligned} \frac{C}{Nk_B} &= (S+s) \left\{ \frac{3\zeta(3/2)}{4\sqrt{2\pi}} \tilde{t}^{1/2} - \tilde{t} \right. \\ &\left. + \frac{15[(S^2 - Ss + s^2)\zeta(5/2) - 4\zeta(1/2)]}{32\sqrt{2\pi}} \tilde{t}^{3/2} \right\} \\ &+ O(\tilde{t}^2), \end{aligned} \quad (4.9)$$

where $\tilde{t} = t/Ss(S+s) = k_B T/JSs(S+s)$. The $O(S^0)$ interactions affect the fourth and higher terms and therefore, whether through the Holstein-Primakoff transformation²⁶ or through the Dyson-Maleev transformation,^{27,36} we reach the same results (4.8) and (4.9).

Numerically solving the thermodynamic Bethe-ansatz integral equations for the spin- $\frac{1}{2}$ ferromagnetic Heisenberg chain, Takahashi and Yamada³¹ obtained

$$\frac{\chi J}{L(g\mu_B)^2} = 0.04167t^{-2} + 0.145t^{-3/2} + 0.17t^{-1}$$

$$+O(t^{-1/2}), \quad (4.10)$$

$$\frac{C}{Lk_B} = 0.7815t^{1/2} - 2.00t + 3.5t^{3/2} + O(t^2), \quad (4.11)$$

where L is the number of spins. When we set S and s both equal to $\frac{1}{2}$, the expressions (4.8) and (4.10) coincide in their leading three terms, while Eqs. (4.9) and (4.11) in their leading two terms. The modified spin-wave calculations are thus reliable and give rigorous information on the low-temperature properties. In the case of arbitrary S and s , the leading three terms of Eq. (4.8) and the leading two terms of Eq. (4.9) coincide with those of the spin- $[Ss(S+s)/2]^{1/3}$ uniform ferromagnetic chain except for a common factor. Considering practical combinations of S and s , we may estimate that $[Ss(S+s)/2]^{1/3} = [1 - (S-s)^2/(S+s)^2]^{1/3}(S+s)/2 \simeq (S+s)/2$. Thus, ferromagnetically coupled alternating spins S and s look like a ferromagnetic assembly of virtual spins $(S+s)/2$ at low temperatures.

It is also interesting to compare the expressions (4.8) and (4.9) with those of ferrimagnetic chains.²⁴ It turns out that the spin- (S,s) ferrimagnetic low-temperature expansions are obtained by replacing J and s by $-J$ and $-s$, respectively, in Eqs. (4.8) and (4.9). In other words, antiferromagnetically coupled alternating spins S and s look like a ferromagnetic assembly of virtual spins $[Ss(S-s)/2]^{1/3} = [(S+s)^2/(S-s)^2 - 1]^{1/3}(S-s)/2$ at low temperatures. The quantity $[Ss(S-s)/2]^{1/3}$ is less intuitive than the corresponding $[Ss(S+s)/2]^{1/3}$ in the case of ferromagnetic coupling, but it results in a realistic spin quantum number $S-s$ when S is equal to $2s$. Equation (4.8) is nothing but the susceptibility of the spin- (S,s) ferrimagnetic chain if we set t for $k_B T/Js(S-s)$ instead of $k_B T/Js(S+s)$. All in all, the low-temperature physics is scaled by $S+s$ in ferromagnetic chains, whereas by $S-s$ in ferrimagnetic chains.

V. CONCLUDING REMARKS

Thermodynamic properties of alternating-spin Heisenberg ferromagnetic chains have been investigated in comparison with heterospin ferrimagnetic and homospin ferromagnetic chains. The magnetic susceptibility is a monotonically decreasing function of temperature re-

gardless of (S,s) and is qualitatively the same as those of uniform ferromagnetic chains. The specific heat qualitatively varies with (S,s) and exhibits a rich structure at intermediate temperatures. It may be double-peaked for $S \gtrsim 3s$ in general.

The low-temperature behavior has been revealed analytically. The thermal quantities are still expanded in powers of $T^{1/2}$ and exhibit ferromagnetic features. The conventional spin-wave theory misunderstands the low-temperature behavior as series of T . The missing terms are reproduced through the modified procedure. Ferromagnetic and ferrimagnetic mixed-spin chains are qualitatively alike at low temperatures. The spin- (S,s) ferromagnetic chain looks like a ferromagnetic assembly of virtual spins $[Ss(S+s)/2]^{1/3} = [1 - (S-s)^2/(S+s)^2]^{1/3}(S+s)/2 \simeq (S+s)/2$, while the spin- (S,s) ferrimagnetic chain behaves like that of virtual spins $[Ss(S-s)/2]^{1/3} = [(S+s)^2/(S-s)^2 - 1]^{1/3}(S-s)/2$. The present findings are really complementary to the sophisticated high-temperature series-expansion calculations.^{20,37}

The existent bimetallic chain ferromagnet $\text{MnNi}(\text{NO}_2)_4(\text{en})_2$ possesses a rather weak intra-chain exchange coupling ($J/k_B \simeq 2\text{K}$), in which the low-temperature thermodynamics revealed here is hard to verify. Nevertheless such a pioneering material must highly motivate further explorations in both chemical and physical fields, as was the case with uniform ferromagnetic chains.^{30,31,32,38,39,40} Besides bimetallic chain compounds, several authors⁴¹ made a novel attempt to design low-dimensional heterospin systems utilizing organic triradicals. Mixed-spin chains contain further interesting topics such as dynamic structure factors of dual aspect⁴² and nuclear spin relaxation through the exchange-scattering-enhanced three-magnon process.⁴³ We hope our study will stimulate further experimental investigations into mixed-spin chain compounds.

Acknowledgments

The authors are grateful to N. Fukushima for valuable comments. This work was supported by the Ministry of Education, Culture, Sports, Science and Technology of Japan, and the Iketani Science and Technology Foundation.

¹ O. Kahn, Y. Pei, and Y. Journaux, in *Inorganic Materials*, edited by D. W. Bruce and D. O'Hare (John Wiley & Sons, New York, 1992), p. 95.

² A. Gleizes and M. Verdaguer, *J. Am. Chem. Soc.* **103**, 7373 (1981); **106**, 3727 (1984).

³ O. Kahn, Y. Pei, M. Verdaguer, J.-P. Renard, and J. Sletten, *J. Am. Chem. Soc.* **110**, 782 (1988); P. J. van Koningsbruggen, O. Kahn, K. Nakatani, Y. Pei, J.-P. Renard, M. Drillon, and P. Legoll, *Inorg. Chem.* **29**, 3325 (1990).

⁴ Y. Pei, M. Verdaguer, O. Kahn, J. Sletten, and J.-P. Renard, *J. Am. Chem. Soc.* **108**, 7428 (1986).

⁵ A. Caneschi, D. Gatteschi, P. Rey, and R. Sessoli, *Inorg. Chem.* **27**, 1756 (1988); A. Caneschi, D. Gatteschi, J.-P. Renard, P. Rey, and R. Sessoli, *ibid.* **28**, 1976 (1989); 2940 (1989).

⁶ A. S. Ovchinnikov, I. G. Bostrem, V. E. Sinityn, A. S. Boyarchenkov, N. V. Baranov, and K. Inoue, *J. Phys.: Condens. Matter* **14**, 8067 (2002).

⁷ M. Drillon, J. C. Gianduzzo, and R. Georges, *Phys. Lett. A* **96**, 413 (1983); M. Drillon, E. Coronado, R. Georges,

- J. C. Gianduzzo, and J. Curely, Phys. Rev. B **40**, 10992 (1989).
- ⁸ M. Verdaguer, A. Gleizes, J.-P. Renard, and J. Seiden, Phys. Rev. B **29**, 5144 (1984).
 - ⁹ A. K. Kolezhuk, H.-J. Mikeska, and S. Yamamoto, Phys. Rev. B **55**, R3336 (1997).
 - ¹⁰ S. Brehmer, H.-J. Mikeska, and S. Yamamoto, J. Phys.: Condens. Matter **9**, 3921 (1997).
 - ¹¹ S. K. Pati, S. Ramasesha, and D. Sen, Phys. Rev. B **55**, 8894 (1997); J. Phys.: Condens. Matter **9**, 8707 (1997).
 - ¹² T. Fukui and N. Kawakami, Phys. Rev. B **55**, R14709 (1997); **56**, 8799 (1997).
 - ¹³ G.-S. Tian, Phys. Rev. B **56**, 5355 (1997).
 - ¹⁴ H. Niggemann, G. Uimin, and J. Zittartz, J. Phys.: Condens. Matter **9**, 9031 (1997); **10**, 5217 (1998).
 - ¹⁵ K. Maisinger, U. Schollwöck, S. Brehmer, H.-J. Mikeska, and S. Yamamoto, Phys. Rev. B **58**, R5908 (1998).
 - ¹⁶ N. B. Ivanov, Phys. Rev. B **62**, 3271 (2000).
 - ¹⁷ O. Kahn, E. Bakalbassis, C. Mathonière, M. Hagiwara, K. Katsumata, and L. Ouahab, Inorg. Chem. **36**, 1530 (1997).
 - ¹⁸ R. Feyerherm, C. Mathonière, and O. Kahn, J. Phys.: Condens. Matter **13**, 2639 (2001).
 - ¹⁹ B. Gillon, C. Mathonière, E. Ruiz, S. Alvarez, A. Cousson, T. M. Rajendiran, and O. Kahn, J. Am. Chem. Soc. **124**, 14433 (2002).
 - ²⁰ N. Fukushima, A. Honecker, S. Wessel, and W. Brenig, Phys. Rev. B **69**, 174430 (2004). N. Fukushima, A. Honecker, S. Wessel, S. Grossjohann, and W. Brenig, Physica B, in press.
 - ²¹ F. Lloret, R. Ruiz, M. Julve, J. Faus, Y. Journaux, I. Castro, and M. Verdaguer, Chem. Mater. **4**, 1150 (1992).
 - ²² J. Larionova, O. Kahn, J. Bartolome, R. Burriel, M. Castro, V. Ksenofontov, and P. Gülich, Chem. Mater. **11**, 3400 (1999).
 - ²³ S. Yamamoto, T. Fukui, K. Maisinger, and U. Schollwöck, J. Phys.: Condens. Matter **10**, 11033 (1998).
 - ²⁴ S. Yamamoto and T. Fukui, Phys. Rev. B **57**, R14008 (1998).
 - ²⁵ F. Bloch, Z. Phys. **61**, 206 (1930); **74**, 295 (1932).
 - ²⁶ T. Holstein and H. Primakoff, Phys. Rev. **58**, 1098 (1940).
 - ²⁷ F. J. Dyson, Phys. Rev. **102**, 1217; 1230 (1956).
 - ²⁸ S. Yamamoto, S. Brehmer, and H.-J. Mikeska, Phys. Rev. B **57**, 13610 (1998).
 - ²⁹ N. B. Ivanov, Phys. Rev. B **57**, R14024 (1998); N. B. Ivanov, J. Richter, and U. Schollwöck, Phys. Rev. B **58**, 14456 (1998).
 - ³⁰ M. Takahashi, Prog. Theor. Phys. Suppl. **87**, 233 (1986).
 - ³¹ M. Takahashi and M. Yamada, J. Phys. Soc. Jpn. **54**, 2808 (1985); M. Yamada and M. Takahashi, J. Phys. Soc. Jpn. **55**, 2024 (1986).
 - ³² M. Takahashi, Phys. Rev. Lett. **58**, 168 (1987).
 - ³³ S. Yamamoto, T. Fukui, and T. Sakai, Eur. Phys. J. B **15**, 211 (2000).
 - ³⁴ S. Yamamoto, Phys. Rev. B **59**, 1024 (1999).
 - ³⁵ T. Nakanishi and S. Yamamoto, Phys. Rev. B **65**, 214418 (2002).
 - ³⁶ S. V. Maleev, Sov. Phys. JETP **6**, 776 (1958).
 - ³⁷ O. Rojas, S. M. de Souza, M. T. Thomaz, J. Math. Phys. **43**, 1390 (2002).
 - ³⁸ C. P. Landee and R. D. Willett, Phys. Rev. Lett. **43**, 463 (1979).
 - ³⁹ P. Schlottmann, Phys. Rev. Lett. **54**, 2131 (1985).
 - ⁴⁰ M. Takahashi, P. Turek, Y. Nakazawa, M. Tamura, K. Nozawa, D. Shiomi, M. Ishikawa, and M. Kinoshita, Phys. Rev. Lett. **67**, 746 (1991).
 - ⁴¹ Y. Hosokoshi, K. Katoh, Y. Nakazawa, H. Nakano, and K. Inoue, J. Am. Chem. Soc. **123**, 7921 (2001).
 - ⁴² S. Yamamoto and T. Sakai, J. Phys. Soc. Jpn. **67**, 3711 (1998).
 - ⁴³ H. Hori and S. Yamamoto, J. Phys. Soc. Jpn. **73**, 1453 (2004); J. Phys.: Condens. Matter **16**, 9023 (2004).




FIG. 1: Single-magnon excitation spectra as the rigorous dispersion relations of the elementary excitations for the spin- (S, s) ferromagnetic Heisenberg chains.




FIG. 2: Modified-spin-wave (MSW) and quantum Monte Carlo (QMC) calculations of the magnetic susceptibility χ as a function of temperature for the spin- (S, s) ferromagnetic Heisenberg chains.




FIG. 3: Modified-spin-wave (MSW) and quantum Monte Carlo (QMC) calculations of the specific heat C as a function of temperature for the spin- (S, s) ferromagnetic Heisenberg chains.




FIG. 4: Modified-spin-wave (MSW) calculations of the specific heat C as a function of temperature for the spin- (S, s) ferromagnetic Heisenberg chains in the cases of $S < 3s$ (a) and $S > 3s$ (b).

